

**ANTO-12-52**

***Oceanibacterium hippocampi* gen. nov., sp. nov., isolated from cutaneous mucus of wild seahorses (*Hippocampus guttulatus*)**

José Luis Balcázar<sup>1,2</sup>, Miquel Planas<sup>1</sup>, José Pintado<sup>1</sup>

<sup>1</sup>Instituto de Investigaciones Marinas, Consejo Superior de Investigaciones Científicas (IIM-CSIC), 36208 Vigo, Spain.

<sup>2</sup>Catalan Institute for Water Research (ICRA), Scientific and Technological Park of the University of Girona, 17003 Girona, Spain.

**Correspondence:** J. L. Balcázar. Tel: +34 972 183 380. Fax: +34 972 183 248.

e-mail: [jlbalcazar@icra.cat](mailto:jlbalcazar@icra.cat)

**Running title:** *Oceanibacterium hippocampi* gen. nov., sp. nov.

**Abstract**

A Gram-negative, aerobic, motile and slightly curved rod-shaped bacterium (BFLP-8<sup>T</sup>) was isolated from cutaneous mucus of wild long-snouted seahorses (*Hippocampus guttulatus*) captured in northwest Spain (Toralla, Galicia). Strain BFLP-8<sup>T</sup> grew at 10–35°C and pH 5–9 (optimally at 25°C and pH 7.0) and with 1–6% (w/v) NaCl (optimally with 2% NaCl). The predominant respiratory quinone (90%) was ubiquinone with ten isoprene units (Q-10) and the major fatty acids identified were C<sub>18:1</sub>ω7c (54.8% of the total), C<sub>19:0</sub> cyclo ω8c (11.6%), C<sub>16:0</sub> (9.5%), C<sub>18:1</sub> 2-OH (7.1%) and C<sub>16:1</sub>ω11c (6.7%). The G+C content of the DNA was 57.8 mol%. Phylogenetic analysis based on 16S rRNA gene sequences showed that strain BFLP-8<sup>T</sup> formed a distinct clade within the family *Sneathiellaceae* but is not specifically associated with any species in the family. On the basis of phenotypic, chemotaxonomic and phylogenetic data, strain BFLP-8<sup>T</sup> represents a novel species within a new genus, for which the name *Oceanibacterium hippocampi* gen. nov., sp. nov. is proposed. The type strain is BFLP-8<sup>T</sup> (= CECT 7691<sup>T</sup> = DSM 23444<sup>T</sup>).

**Keywords:** *Oceanibacterium hippocampi*; polyphasic taxonomic analysis; seahorses

## Introduction

The class *Alphaproteobacteria* comprises a large group of Gram-negative bacteria within the phylum *Proteobacteria* and is currently divided into ten orders: *Caulobacterales*, *Kiloniellales*, *Kordiimonadales*, *Parvularculales*, *Rhizobiales*, *Rhodobacterales*, *Rhodospirillales*, *Rickettsiales*, *Sneathiellales* and *Sphingomonadales* (www.bacterio.cict.fr). These bacteria show a wide range of morphological, physiological and genetic characteristics (Batut *et al.* 2004).

The family *Sneathiellaceae* in the order *Sneathiellales* was proposed by Kurahashi *et al.* (2008) based on phylogenetic analysis of 16S rRNA gene sequences. This family is currently represented by one genus with two species: *Sneathiella chinensis*, isolated from coastal sediments in China (Jordan *et al.* 2007) and *Sneathiella glossodoripedis*, isolated from the foot epidermis of a nudibranch in Japan (Kurahashi *et al.* 2008). In the present study, we describe the physiological, chemotaxonomic and phylogenetic characteristics of a novel strain, BFLP-8<sup>T</sup>, belonging to a new genus of the family *Sneathiellaceae*.

## Materials and Methods

### Strain isolation

During the characterization of organisms isolated from cutaneous mucus of wild long-snouted seahorses (*Hippocampus guttulatus*), strain BFLP-8<sup>T</sup> was isolated on marine agar (Difco, Becton Dickinson, Heidelberg, Germany) at 20°C. The organism was subcultured on the same medium at 25°C for 3 days. Stock cultures were stored at –80 °C in marine broth (Difco) with 30% (v/v) glycerol.

### Phenotypic tests

Gram reaction was determined using the non-staining (KOH) method as described by Buck (1982). Cell morphology and motility were studied using phase-contrast

microscopy and electron microscopy as previously described by Herrera *et al.* (2007). NaCl growth tolerance and requirements were investigated by using nutrient broth [0.5% peptone from casein, 0.3% meat extract, 0.3% yeast extract, adjusted to pH 7.2] supplemented with various concentrations of NaCl (0–15% at intervals of 1%). The pH range for growth was determined in marine broth that was adjusted to various pH values with acetic acid-sodium acetate (pH 4.0-4.5, 100 mM), MES (pH 5.0-6.0, 50 mM), MOPS (pH 6.5, 50 mM), Tris (pH 7.0-9.0, 50 mM) or CHES (pH 9.5-10.0, 50 mM) buffers. Anaerobic growth was assessed at 25 °C in anaerobic chambers with an H<sub>2</sub>/CO<sub>2</sub> atmosphere (bioMérieux).

Catalase activity was determined by assessing bubble production in 3% (v/v) H<sub>2</sub>O<sub>2</sub>; oxidase activity was determined using 1% (w/v) tetramethyl-*p*-phenylenediamine as described by Lim *et al.* (2008). Some physiological characteristics were performed using API 20NE and API ZYM (bioMérieux). Cells for inoculation of the strips were grown for 3 days at 25°C on marine agar and the results were visually interpreted according to the manufacturer's instructions.

For base composition analysis, DNA was prepared according to Chun & Goodfellow (1995) and the G+C content was determined using the thermal denaturation method (Mandel & Marmur, 1968). DNA from *Escherichia coli* ATCC 11775<sup>T</sup> was used as the reference for determination of the thermal-melting profile ( $T_m$ ).

#### **Chemotaxonomic analyses**

Whole-cell fatty acids from the isolate were extracted from biomass grown on marine agar and were analysed according to the standard protocol of the Sherlock Microbial Identification System (MIDI version 4.5). Analysis of respiratory quinones was carried out by the DSMZ Identification Service and Dr. Brian Tindall (DSMZ, Braunschweig, Germany).

## Phylogenetic analysis

Extraction and amplification of genomic DNA for 16S rRNA sequence analysis were carried out as described previously (Balcázar *et al.* 2010). The sequences obtained were compared against the sequences available in the GenBank, EMBL and DDBJ databases obtained from the National Center for Biotechnology Information using the BLASTN (Altschul *et al.* 1990). Phylogenetic analysis was performed using the software MEGA version 5.0 (Tamura *et al.* 2011) after multiple alignments of data by CLUSTAL W (Larkin *et al.* 2007). Distances (distance options according to the Kimura two-parameter model) and clustering with the neighbour-joining and maximum-parsimony methods were determined using bootstrap values based on 1000 replications.

## Results and Discussion

Cells of strain BFLP-8<sup>T</sup> were slightly curved rods (Fig. 1), Gram-negative, oxidase- and catalase-positive, motile and aerobic. In addition, strain BFLP-8<sup>T</sup> could be differentiated from its closest phylogenetic neighbours (see below) on the basis of some biochemical properties such as nitrate reduction, arginine dihydrolase and urease activities. Other physiological characteristics of strain BFLP-8<sup>T</sup> are shown in Table 1 and also in the species description.

The DNA G+C content was calculated to be 57.8 mol%. This value corresponds closely to the range (56.9–57.1 mol%) reported for recognized members of the family *Sneathiellaceae* (Jordan *et al.* 2007; Kurahashi *et al.* 2008).

The predominant respiratory quinone of strain BFLP-8<sup>T</sup> was Q-10 (90%), with Q-9 present in minor amounts (10%). Q-10 has been reported as the predominant quinone in many members of the *Alphaproteobacteria* (Wagner-Döbler *et al.* 2004), including species of the genus *Sneathiella* (Kurahashi *et al.* 2008). In addition, the fatty acid profile was composed of C<sub>18:1</sub>ω7c (54.8%), C<sub>19:0</sub> cyclo ω8c (11.6%), C<sub>16:0</sub> (9.5%), C<sub>18:1</sub>

2-OH (7.1%), C<sub>16:1</sub>ω11c (6.7%), 11-methyl-C<sub>18:1</sub> ω7c (2.6%), C<sub>18:1</sub> ω9c (2.2%), C<sub>18:0</sub> (2.0%) and C<sub>18:0</sub> 3-OH (2.0%). Fatty acids C<sub>16:0</sub> and C<sub>18:1</sub>ω7c are typically the major fatty acids found in members of the family *Sneathiellaceae* (Kurahashi *et al.* 2008).

The GenBank/EMBL/DDBJ accession number for the 16S rRNA gene sequence of strain BFLP-8<sup>T</sup> is FN687912. 16S rRNA gene sequence similarity calculations after a neighbour-joining analysis indicated that strain BFLP-8<sup>T</sup> grouped most closely with uncultured bacterial clones (GenBank accession numbers FJ202709 and FJ205319, 92.6 and 92.4% sequence identity). The closest cultured relatives of strain BFLP-8<sup>T</sup> were *Oceanibaculum pacificum* MC2UP-L3<sup>T</sup> (91.5%), *Sneathiella glossodoripedis* IAM 15419<sup>T</sup> (91.1%), *Sneathiella chinensis* LMG 23452<sup>T</sup> (90.9%) and *Oceanibaculum indicum* P24<sup>T</sup> (90.5%). Similar results were obtained with the maximum-parsimony algorithm (Fig. S1). The phylogenetic tree suggested that the novel strain forms a distinct clade within the family *Sneathiellaceae* but is not specifically associated with any species in the family (Fig. 2). Phylogenetic analyses and the low levels of 16S rRNA gene sequence similarities between the novel strain and other members of the family *Sneathiellaceae* indicate that strain BFLP-8<sup>T</sup> represent a new genus in the family. Therefore, the phenotypic and genotypic properties of strain BFLP-8<sup>T</sup> support its description as a novel species within a new genus, for which the name *Oceanibacterium hippocampi* gen. nov., sp. nov. is proposed.

#### Description of *Oceanibacterium* gen. nov.

*Oceanibacterium* (L. n. *oceanus*, the ocean; L. neut. n. *bacterium*, rod; N.L. neut. n. *Oceanibacterium*, rod-shaped bacterium from the ocean [sea water]).

Gram-negative, motile, aerobic, non-spore-forming and rod-shaped. Cells grow at temperatures in the range 10–35 °C. NaCl is required for growth. Positive for oxidase and catalase activities. Nitrate is reduced to nitrite. The ubiquinone system is Q-10 and

the major fatty acids (>6%) are C<sub>18:1</sub>ω7c, C<sub>19:0</sub> cyclo ω8c, C<sub>16:0</sub>, C<sub>18:1</sub> 2-OH and C<sub>16:1</sub>ω11c. The type species is *Oceanibacterium hippocampi*.

**Description of *Oceanibacterium hippocampi* sp. nov.**

*Oceanibacterium hippocampi* (hip.po.cam'pi. L. gen. n. *hippocampi*, of the seahorse, isolated from *Hippocampus guttulatus*).

Displays the following properties in addition to those given in the genus description.

Cells are slightly curved rod-shaped (0.4 × 1.0–1.6 μm). Optimum growth temperature is 25°C. No growth occurs below 10°C or above 35°C. Growth occurs at pH 5.0–9.0, but not below pH 4.5 or above pH 9.5. Growth occurs at NaCl concentrations between 1.0 and 6.0% (w/v) but not without NaCl or in the presence of >7.0% NaCl (w/v).

Positive for arginine dihydrolase, urease and assimilation of adipate and malate.

Negative for indole production, glucose fermentation, aesculin, gelatine hydrolysis, β-

galactosidase and assimilation of D-glucose, L-arabinose, D-mannose, D-mannitol, N-

acetyl-D-glucosamine, D-maltose, potassium gluconate, caprate, citrate and phenyl-

acetate. API ZYM tests show activities for alkaline phosphatase, esterase (C4), esterase

lipase (C8), lipase (C14), leucine arylamidase, valine arylamidase, cystine arylamidase,

acid phosphatase and naphthol-AS-BI-phosphohydrolase but not for trypsin, α-

chymotrypsin, α-galactosidase, β-galactosidase, β-glucuronidase, α-glucosidase, β-

glucosidase, α-mannosidase and α-fucosidase. The major fatty acids are C<sub>16:0</sub>, C<sub>16:1</sub>ω11c,

C<sub>18:1</sub>ω7c, C<sub>19:0</sub> cyclo ω8c and C<sub>18:1</sub> 2-OH; smaller amounts of C<sub>18:0</sub>, C<sub>18:1</sub>ω9c, 11-

methyl-C<sub>18:1</sub>ω7c and C<sub>18:0</sub> 3-OH are present. The G+C content of the type strain is 57.8

mol% (T<sub>m</sub>).

The GenBank/EMBL/DDBJ accession number for the 16S rRNA gene sequence of strain BFLP-8<sup>T</sup> is FN687912.

The type strain, BFLP-8<sup>T</sup> (= CECT 7691<sup>T</sup> = DSM 23444<sup>T</sup>), was isolated from cutaneous mucus of wild long-snouted seahorses (*Hippocampus guttulatus*).

**Acknowledgements**

This study was financed by the Spanish Ministry of Science and Technology (Projects CGL2005-05927-C03-01 and CGL2009-08386) and by the Regional Government Xunta de Galicia (09MDS022402PR). J.L.B. was supported by a postdoctoral I3P contract from the Spanish National Research Council (CSIC), co-financed by the European Social Fund. We thank P. Quintas, A. Chamorro, M. Cueto, S. Otero and P. Ruiz for skilful technical assistance.

**References**

- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. *J Mol Biol* 215:403–410.
- Balcázar JL, Pintado J, Planas M (2010) *Bacillus galliciensis* sp. nov., isolated from faeces of wild seahorses (*Hippocampus guttulatus*). *Int J Syst Evol Microbiol* 60:892–895.
- Batut J, Andersson SGE, O’Callaghan D (2004) The evolution of chronic infection strategies in the alpha-proteobacteria. *Nat Rev Microbiol* 2:933–945.
- Buck JD (1982) Nonstaining (KOH) method for determination of Gram reactions of marine bacteria. *Appl Environ Microbiol* 44:992–993.
- Chun J, Goodfellow M (1995) A phylogenetic analysis of the genus *Nocardia* with 16S rRNA gene sequences. *Int J Syst Bacteriol* 45:240–245.
- Dong C, Lai Q, Chen L, Sun F, Shao Z, Yu Z (2010) *Oceanibaculum pacificum* sp. nov., isolated from hydrothermal field sediment of the south-west Pacific Ocean. *Int J Syst Evol Microbiol* 60:219–222.



- 200 Herrera JJR, Cabo ML, González A, Pazos I, Pastoriza L (2007) Adhesion and  
201 detachment kinetics of several strains of *Staphylococcus aureus* subsp. *aureus* under  
202 three different experimental conditions. Food Microbiol 24:585–591.
- 203 Jordan EM, Thompson FL, Zhang XH, Li Y, Vancanneyt M, Kroppenstedt RM, Priest  
204 FG, Austin B (2007) *Sneathiella chinensis* gen. nov., sp nov., a novel marine  
205 alphaproteobacterium isolated from coastal sediment in Qingdao, China. Int J Syst Evol  
206 Microbiol 57:114–121.
- 207 Kurahashi M, Fukunaga Y, Harayama S, Yokota A (2008) *Sneathiella glossodoripedis*  
208 sp. nov., a marine alphaproteobacterium isolated from the nudibranch *Glossodoris*  
209 *cincta*, and proposal of *Sneathiellales* ord. nov. and *Sneathiellaceae* fam. nov. Int J Syst  
210 Evol Microbiol 58:548–552.
- 211 Lai Q, Yuan J, Wu C, Shao Z (2009) *Oceanibaculum indicum* gen. nov., sp. nov.,  
212 isolated from deep seawater of the Indian Ocean. Int J Syst Evol Microbiol 59:1733–  
213 1737.
- 214 Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H,  
215 Valentin F, Wallace IM, Wilm A, Lopez R, Thompson JD, Gibson TJ, Higgins DG  
216 (2007) Clustal W and Clustal X version 2.0. Bioinformatics 23:2947–2948.
- 217 Lim JM, Jeon CO, Jang HH, Park DJ, Shin YK, Yeo SH, Kim CJ (2008) *Albimonas*  
218 *donghaensis* gen. nov., sp. nov., a non-photosynthetic member of the class  
219 *Alphaproteobacteria* isolated from seawater. Int J Syst Evol Microbiol 58:282–285.
- 220 Mandel M, Marmur J (1968) Use of ultraviolet absorbance temperature profile for  
221 determining the guanine plus cytosine content of DNA. Methods Enzymol 12B:195–  
222 206.

223 Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S (2011) MEGA5:  
224 Molecular Evolutionary Genetics Analysis using Maximum Likelihood, Evolutionary  
225 Distance, and Maximum Parsimony Methods. Mol Biol Evol 28:2731–2739.

226 Wagner-Döbler I, Rheims H, Felske A, El-Ghezal A, Flade-Schröder D, Laatsch H,  
227 Lang S, Pukall R, Tindall BJ (2004) *Oceanibulbus indolifex* gen. nov., sp. nov., a North  
228 Sea alphaproteobacterium that produces bioactive metabolites. Int J Syst Evol Microbiol  
229 54:1177–1184.

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244


245 **Table 1.** Differential characteristics of strain BFLP-8<sup>T</sup> and some related species

Characteristic	1	2	3	4	5
Oxidase	+	+	+	+	–
Growth without NaCl	–	–	+	+	+
Growth at 45 °C	–	+	–	+	–
Nitrate reduction	+	+	+	–	+
Arginine dihydrolase	+	+	–	+	–
Urease	+	–	+	+	–
Citrate utilization	–	+	–	–	–

246  
247 Strains: 1, *Oceanibacterium hippocampi* gen. nov., sp. nov. BFLP-8<sup>T</sup>; 2, *Sneathiella*  
248 *glossodoripedis* MKT133<sup>T</sup>; 3, *Sneathiella chinensis* LMG 23452<sup>T</sup>; 4, *Oceanibaculum*  
249 *pacificum* LMC2up-L3<sup>T</sup>; 5, *Oceanibaculum indicum* P24<sup>T</sup>. +, positive; –, negative. Data  
250 for 2-5 taken from Kurahashi *et al.* (2008), Jordan *et al.* (2007), Dong *et al.* (2010) and  
251 Lai *et al.* (2009), respectively.

252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279

280



281

282

283

284

**Fig. 1.** Scanning electron micrograph of strain BFLP-8<sup>T</sup> showing a slightly curved, rod-

285

shaped morphology ( $0.4 \times 1.0\text{--}1.6\ \mu\text{m}$ ). Bar,  $0.5\ \mu\text{m}$ .

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304


305

306


307

308

309



**Fig. 2.** Phylogenetic analysis of strain BFLP-8<sup>T</sup> and representatives of the *Alphaproteobacteria* based on 16S rRNA gene sequences (GenBank/EMBL/DDBJ accession numbers in parentheses), constructed by the neighbour-joining method. Bootstrap percentages (>50%) based on 1000 replications are shown at branch nodes. *Escherichia coli* ATCC 11775<sup>T</sup> was used as an outgroup. Bar, 0.02 substitutions per nucleotide position.



**Supplementary Fig. S1.** Phylogenetic analysis of strain BFLP-8<sup>T</sup> and representatives of the *Alphaproteobacteria* based on 16S rRNA gene sequences (GenBank/EMBL/DBJ accession numbers in parentheses), constructed after multiple alignment of data by CLUSTAL W (Larkin *et al.* 2007). Distances were calculated and clustering with the maximum-parsimony method was performed by using the software package MEGA version 5.0 (Tamura *et al.* 2011). Bootstrap values based on 1000 replications are listed as percentages at branching points. *Escherichia coli* ATCC 11775<sup>T</sup> was used as an outgroup. Bar, 10 substitutions per nucleotide position.

## References

- Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, Valentin F, Wallace IM, Wilm A, Lopez R, Thompson JD, Gibson TJ, Higgins DG (2007) Clustal W and Clustal X version 2.0. *Bioinformatics* 23:2947–2948.
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S (2011) MEGA5: Molecular Evolutionary Genetics Analysis using Maximum Likelihood, Evolutionary Distance, and Maximum Parsimony Methods. *Mol Biol Evol* 28:2731–2739.